

## Clinician Management Resource for individuals with a germline predisposition to myelodysplastic syndrome (MDS) or acute myeloid leukemia (AML)

This overview of clinical management guidelines is based on this patient's positive test result. Unless otherwise stated, medical management guidelines used here are limited to those published in the Nordic Guidelines for Germline Predisposition to Myeloid Neoplasms in Adults<sup>1</sup> since U.S. consensus practice guidelines have not been developed. Please consult the referenced guideline for complete details and further information.

Clinical correlation with the patient's past medical history, treatments, surgeries, and family history may lead to changes in clinical management decisions; therefore, other management recommendations may be considered. Genetic testing results and published guidelines help inform medical management decisions but do not constitute formal recommendations. Discussions of medical management decisions and individualized treatment plans should be made in consultation between each patient and his or her healthcare provider and may change.

SCREENING/SURGICAL CONSIDERATIONS <sup>1</sup>	AGE TO START	FREQUENCY	
Myelodysplastic syndrome (MDS) and Acute Myeloid Leukemia (AML)*			
Complete blood count (CBC) with manual differential	Baseline	Every 6 months	
Bone marrow aspirate/biopsy with cytogenetic analysis	Baseline	Repeat only if CBC changes	
Testing for somatic gene mutations using a next generation sequencing myeloid gene panel with high coverage and reading depth	Baseline (bone marrow)	Annually (blood)	
Management and surveillance of other organ disfunction			
Referral to a hereditary cancer clinic or to relevant medical specialists to ensure screening for solid tumors and organ disfunction	Individualized, depending on the underlying condition	Individualized, depending on the underlying condition	
Other			
Referral to genetic counseling when family planning is relevant, preferably before pregnancy	Individualized	Individualized	
Consideration of allogenic hematopoietic stem cell transplant (allo-HSCT)**	Individualized	Individualized	

\* Very little evidence-based data exist on the efficacy and benefit of surveillance in individuals with germline predisposition to MDS and AML, and published recommendations for surveillance are based on expert opinion.<sup>1</sup>

\*\* All patients of a suitable age who have developed myeloid neoplasms on the basis of a genetic predisposition, except those diagnosed with AML associated with germline variants in CEBPA, are potential candidates for allo-HSCT. Each case should be referred for discussion with an expert transplantation panel that may include international specialists in the field.

 Baliakas P, Tesi B, Wartiovaara-Kautto U, Stray-Pedersen A, FriisL S, Dybedal I, Hovland R, Jahnukainen K, Raaschou-Jensen K, Ljungman P, Rustad CF, Lautrup CK, Kilpivaara O, Kittang AO, Grønbæk K, Cammenga J, Hellström-Lindberg E, Andersen MK. Nordic guidelines for germline predisposition to myeloidneoplasms in adults: Recommendations for genetic diagnosis, clinical management and follow-up. HemaSphere, 2019;3:6. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6924562/



## Understanding Your Positive ETV6 Genetic Test Result


INFORMATION FOR PATIENTS WITH A PATHOGENIC OR LIKELY PATHOGENIC VARIANT

## 5 Things To Know


1	Result	Your testing shows that you have a pathogenic or likely pathogenic variant in the <i>ETV6</i> gene. A pathogenic or likely pathogenic <i>ETV6</i> variant may be in a person's body from birth (germline), or may have occurred at some point during a person's lifetime (somatic).
2	<i>ETV6</i> -related thrombocytopenia	People with germline pathogenic or likely pathogenic <i>ETV6</i> variants have <i>ETV6</i> -related thrombocytopenia, a condition that causes low platelets in the blood and can result in mild to moderate bleeding tendency and/or easy bruising.
3	Cancer risks	People with germline pathogenic or likely pathogenic <i>ETV6</i> variants have an increased chance to develop blood cancers. The most common associated cancer is B-cell acute lymphoblastic leukemia (B-ALL), though some people may develop other types of leukemia, lymphoma, or a blood disorder known as myelodysplastic syndrome (MDS). Somatic pathogenic or likely pathogenic <i>ETV6</i> variants do not run in families and cannot be passed on to children; however, they may occur because of an active blood disorder such as leukemia. It is important to discuss your result in detail with your healthcare provider.
4	What you can do	Risk management decisions are very personal. There are options to detect cancer early or lower the risk to develop cancer. It is important to discuss these options with your healthcare provider and decide on a plan that works for you.
5	Family	Family members may also be at risk – they can be tested for the pathogenic or likely pathogenic <i>ETV6</i> variant that was found in you. It is recommended that you share this information with your family members so they can learn more and discuss with their healthcare providers.

## ETV6 Lifetime Cancer Risks\*

For people with germline pathogenic or likely pathogenic variants



\* Because risk estimates vary in different studies, only approximate risks are given. Cancer risks will differ based on individual and family history.



ETV6 in the Family

There is a 50/50 random chance to pass on a germline pathogenic or likely pathogenic *ETV6* variant to each of

Has a germline pathogenic or likely pathogenic *ETV6* variant

- No germline pathogenic or likely pathogenic *ETV6* variants
- American Cancer Society cancer.org
  Genetic Alliance geneticalliance.org
  National Organization for Rare Diseases (NORD) rarediseases.org
  National Society of Genetic Counselors nsgc.org
  Canadian Society of Genetic Counsellors cagc-accg.ca

Please discuss this information with your healthcare provider. The cancer genetics field is continuously evolving, so updates related to your *ETV6* result, medical recommendations, and/or potential treatments may be available over time. This information is not meant to replace a discussion with a healthcare provider, and should not be considered or interpreted as medical advice.